

УДК 517.984.52

\*G. Dildabek, \*\*A. Tengayeva

\*Al-Farabi Kazakh National University, Respublika Kazakhstan, Almaty

\*\*Kazakh National Agrarian University, Respublika Kazakhstan, Almaty

E-mail: \*dildabek.g@gmail.com, \*\*aijan0973@mail.ru

### Constructing a basis from systems of eigenfunctions of one not strengthened regular boundary value problem

In the present work we investigate a nonlocal boundary value spectral problem for an ordinary differential equation in an interval. Such problems arise in solving the nonlocal boundary value for partial equations by the Fourier method of variable separation. For example, they arise in solving nonstationary problems of diffusion with boundary conditions of Samarskii Ionkin type. Or they arise in solving problems with stationary diffusion with opposite flows on a part of the interval. The boundary conditions of this problem are regular but not strengthened regular. The principal difference of this problem is: the system of eigenfunctions is complete but not forming a basis. Therefore the direct applying of the Fourier method is impossible. Based on these eigenfunctions there is constructed a special system of functions that already forms the basis. However the obtained system is not already the system of the eigenfunctions of the problem. In the paper we demonstrate how this new system of functions can be used for solving a nonlocal boundary value equation on the example of the Laplace equation.

**Key words:** nonlocal boundary conditions; regular but not strengthened regular conditions; basis; eigenfunctions; biorthogonal system.

Г. Дилдабек, А.А. Тенгаева

### Построение базиса из системы собственных функций одной неусиленно регулярной краевой задачи

В настоящей работе мы исследуем нелокальную граничную спектральную задачу для обыкновенного дифференциального уравнения на отрезке. Задачи подобного вида возникают при решении методом разделения переменных Фурье нелокальной краевой задачи для уравнений в частных производных. Например, при решении нестационарных задач диффузии с краевыми условиями типа Самарского Ионкина. Или при решении задач стационарной диффузии с противоположными потоками на части границы. Граничные условия этой задачи являются регулярными, но не усиленно регулярными. Принципиальным отличием этой задачи является то, что система собственных функций является полной и минимальной, но не образует базиса. Поэтому прямое применение метода Фурье оказывается невозможным. Основываясь на этих собственных функциях в работе построена специальная система функций, которые уже образуют базис. Однако полученная система уже не является системой собственных функций задачи. В работе демонстрируется, как эта новая система функций может быть использована для решения нелокальной краевой задачи на примере уравнения Лапласа.

**Ключевые слова:** нелокальные граничные условия; регулярные, но не усиленно регулярные краевые условия; базис; собственные функции; биортогональная система.

Г. Дилдабек, А.А. Тенгаева

### Бір қатаң емес регуляр шеттік есептің меншікті функциялар жүйесінен базис құру

Бұл жұмыста кесіндідегі жәй дифференциалдық теңдеу үшін бейлокал шекаралық спектралды есебі зерттелінеді. Мұндай есептер дербес туындылы дифференциалдық теңдеулер үшін бейлокал шеттік есептерді Фурьенің айнмалыны ажырату әдісімен шешу кезінде пайда болады. Мысалы, Самарский Ионкин тектес шеттік шартпен берілген диффузияның бейстационар есептерін шешу кезінде. Немесе шекараның бөлігінде қарама қарсы ағынмен берілген диффузияның стационар есептерін шешу кезінде.

Бұл есептің шекаралық шарты регуляр, бірақ қатаң емес регуляр болып табылады. Бұл есептің айрықша ерекшелігі, оның меншікті функциялар жүйесі базис құрамайды. Сондықтан Фурье әдісін тікелей қолдану мүмкін емес. Осы меншікті функцияларды негізге ала отырып, базис құрайтын арнайы функциялар жүйесі құрылған. Бірақ алынған жүйе берілген есептің меншікті функциялар жүйесі болмайды. Жұмыста осы алынған жаңа функциялар жүйесінің қолданылуына мысал ретінде Лаплас теңдеуі үшін бейлокал шеттік есепті шешілуі көрсетілген.

**Түйін сөздер:** бейлокал шекаралық шарт, регуляр, бірақ қатаң емес регуляр шеттік шарт, меншікті функциялар, биортогонал жүйе.

## Introduction

Investigations on spectral theory of ordinary differential operators begun from classical papers of J. Liouville and Sh. Sturm. Fundamental works in the spectral theory of differential operators were the papers by Birkhoff of 1908, where he introduced regular boundary conditions for the first time. The theory was significantly developed by Tamarkin and Stone. These works led to a new wide scientific direction having an enormous literature. We refer to [1, 2] for the extensive bibliography and the obtained results.

Despite the apparent simplicity, the spectral theory of ordinary differential operators is far from complete. This applies even to the case of a second-order operator

$$Lu = u''(x) + q(x)u$$

on the finite interval  $x \in (a, b)$  which is called Sturm-Liouville operator. Brief survey of results in the spectral theory of the Sturm-Liouville operator is given in the recent paper by Makin [3].

It is known that boundary conditions can be divided into three classes [4]:

- strengthened regular conditions;
- regular but not strengthened regular conditions;
- irregular conditions.

If the boundary conditions are strengthened regular then the system of root functions forms a Riesz basis in  $L_2(a, b)$ . This statement was proved in [5, 6] and [7, Chapter XIX].

In the other cases the basis property of the systems of root functions is not guaranteed. The final definition of classes of the boundary conditions for an operator of second order when the system of eigen- and associated functions forms the basis, was given in [8].

In the present work we consider one model spectral problem for an operator of multiple differentiation. Boundary conditions of the problem are regular but not strengthened regular. The system of eigenfunctions of the problem is complete, minimal, almost normed, but does not form a basis in  $L_2$ . On the basis of these eigenfunctions we construct a special system having basis property in  $L_2$ .

## Statement of the problem

Consider the spectral problem

$$-u''(x) = \lambda u(x), \quad 0 < x < \pi; \tag{1}$$

$$u(0) = 0, \quad u'(0) + u'(\pi) + \alpha u(\pi) = 0,$$

where  $\alpha > 0$  is a fixed parameter.

This problem arises while solving a nonlocal boundary value problem for the Laplace equation by the method of separation of variables. Let  $D = \{(r, \theta) : 0 < r < 1, 0 < \theta < \pi\}$  be a half-disc. Our goal is to find a function  $u(r, \varphi) \in C^0(\bar{D}) \cap C^2(D)$  satisfying in  $D$  the equation

$$\Delta U = 0 \tag{2}$$

with the boundary conditions

$$U(1, \theta) = f(\theta), \quad 0 \leq \theta \leq \pi, \tag{3}$$

$$u(r, 0) = 0, \quad r \in [0, 1], \tag{4}$$

$$\frac{\partial U}{\partial \theta}(r, 0) + \frac{\partial U}{\partial \theta}(r, \pi) + \alpha U(r, \pi) = 0, \quad r \in (0, 1). \tag{5}$$

The difference of this problem is the impossibility of direct applying of the Fourier method (separation of variables). Because the corresponding spectral problem for the ordinary differential equation has the system of eigenfunctions not forming a basis. For  $\alpha = 0$  the problem (2) - (5) was considered in [9].

One method of constructing the basis, based on the system of eigenfunctions of the problem

$$-\vartheta''(x) = \lambda \vartheta(x), \quad 0 < x < \pi;$$

$$\vartheta(0) = 0, \quad \vartheta'(0) = \vartheta'(\pi) + \alpha \vartheta(\pi)$$

was suggested in [11].

The boundary conditions of this problem are regular but not strengthened regular conditions. And the system of its eigenfunctions does not form the basis. But a special system of functions built with help of these eigenfunctions will form the basis. And this fact is applied for the solution of a nonlocal initial-boundary problem for the heat equation.

The goal of the present work is to construct the basis from the system of the eigenfunctions of the problem (1).

## Preliminaries

Let us present briefly the main definitions and facts which will be used in what follows. Let  $B$  be a Banach space with the norm  $\|\cdot\|_B$ , and let  $B^*$  be its dual with the norm  $\|\cdot\|_{B^*}$ .

A system of elements  $\{\varphi_k\}_{k=0}^{\infty}$  is said to be closed in  $B$  if the linear span of this system is everywhere dense in  $B$ ; that is, any element of the space  $B$  can be approximated by a linear combination of elements of this system with any accuracy in the norm of the space  $B$ .

A system of elements  $\{\varphi_k\}_{k=0}^{\infty}$  is said to be minimal in  $B$  if none of its elements belongs to the closure of the linear span of the other elements of this system.

It is well known that a system  $\{\varphi_k\}_{k=0}^{\infty}$  is minimal if and only if there exists a biorthogonal system dual to it, that is, a system of linear functionals  $\{\psi_k\}_{k=0}^{\infty}$  from  $B^*$  such that

$$(\varphi_k, \psi_j) = \delta_{k,j}$$

for all  $k, j \in \mathbb{N}$ . Moreover, if the initial system is simultaneously closed and minimal in  $B$ , then the system biorthogonally dual to it is uniquely defined.

We say that a system  $\{\varphi_k\}_{k=0}^{\infty}$  is uniformly minimal in  $B$ , if there exists  $\gamma > 0$  such that for all  $k \in \mathbb{N}$ ,

$$\text{dist}\{\varphi_k, B_k\} \geq \gamma \|\varphi_k\|_B,$$

where  $B_k$  is the closure of the linear span of all elements  $\varphi_l$  with serial numbers  $l \neq k$ .

It is also well known that a closed and minimal system  $\{\varphi_k\}_{k=0}^{\infty}$  is uniformly minimal in  $B$  if and only if:

$$\sup_{k \in \mathbb{N}} \|\varphi_k\|_B \|\psi_k\|_{B^*} < \infty.$$

A system  $\{\varphi_k\}_{k=0}^{\infty}$  forms a basis of the space  $B$  if, for any element  $f \in B$ , there exists a unique expansion of it in the elements of the system, that is, the series  $\sum_{k=0}^{\infty} f_k \varphi_k$  convergent to  $f$  in the norm of the space  $B$ .

Any basis is a closed and minimal system in  $B$ , and, therefore, we can uniquely find its biorthogonal dual system  $\{\psi_k\}_{k=0}^{\infty}$ , and hence the expansion of any element of  $f$  with respect to the basis  $\{\varphi_k\}_{k=0}^{\infty}$  coincides with its biorthogonal expansion, that is,  $f_k = (f, \psi_k)$  for all  $k \in \mathbb{N}$ .

### On eigenvalues and eigenfunctions of the problem

In a whole the constructing eigenvalues and eigenfunctions of the problem (1) is a simple task. Therefore we omit some details of the calculations and present the main facts which we will use further.

We look for eigenvalues of the problem. Note that  $\lambda = 0$  is not an eigenvalue, since problem (1) for this value of  $\lambda$  has only the trivial solution.

Let  $\lambda \neq 0$ . The eigenfunction should have the form  $u(x) = \sin(\sqrt{\lambda}x)$ . By taking into account the nonlocal boundary condition, we obtain two equations

$$\cos\left(\frac{\sqrt{\lambda}\pi}{2}\right) = 0, \quad \cot\left(\frac{\sqrt{\lambda}\pi}{2}\right) = -\frac{\alpha}{\sqrt{\lambda}}.$$

Solutions of the first equation form a series of eigenvalues and eigenfunctions of the problem (1) of the form

$$\lambda_k^{(1)} = (2k+1)^2, \quad u_{k1}(x) = \sin((2k+1)x), \quad k = 0, 1, 2, \dots$$

The second equations can be represented as

$$\cot(\beta\pi) = -\frac{\alpha}{2\beta}, \quad \beta = \frac{\sqrt{\lambda}}{2}.$$

By  $\beta_k$  denote roots of this equation. It is easy to show that they satisfy the inequalities  $2k+1 < 2\beta_k < 2k+2$ ,  $k = 0, 1, 2, \dots$ , and two-side estimates are carried out for  $\delta_k = \beta_k - k - 1/2$  where  $k$  is large enough

$$\frac{\alpha}{\pi(2k+1)} \left(1 - \frac{1}{2k+1}\right) < \delta_k = \alpha \left|O\left(\frac{1}{k}\right)\right| < \frac{\alpha}{\pi(2k+1)}. \quad (6)$$

Consequently there exists a second series of eigenvalues and eigenfunctions of the form

$$\lambda_k^{(2)} = (2\beta_k)^2, \quad u_{k2}(x) = \sin(2\beta_k x), \quad k = 0, 1, 2, \dots$$

**Lemma 1.** *The system of eigenfunctions  $\{u_{k1}, u_{k2}\}_{k=0}^{\infty}$  of the problem (1) is complete and minimal, almost normed but does not form even an ordinary basis in  $L_2(0, \pi)$ .*

**Proof.** The completeness and minimality of the system follow from the regularity of boundary conditions of the spectral problem (1). The limitation of norms is easily checked by direct calculation. However the properties of the completeness and minimality are not enough for the basis property.

Really, consider scalar multiplications of pairs of eigenfunctions  $(u_{k1}, u_{k2})$ . By direct calculation, we find

$$(u_{k1}, u_{k2}) = \int_0^\pi \sin((2k+1)t) \sin(2\beta_k t) dt = \frac{\pi \sin(2\delta_k \pi)}{2} \frac{2k+1}{2k+1+\delta_k}.$$

Taking into account that  $\|u_{k1}\| = \sqrt{\pi/2}$ , and  $\lim_{k \rightarrow \infty} \|u_{k2}\| = \sqrt{\pi/2}$ , we get that the angle between the normed eigenvectors tends to zero:

$$\lim_{k \rightarrow \infty} \left( \frac{u_{k1}}{\|u_{k1}\|}, \frac{u_{k2}}{\|u_{k2}\|} \right)_{L_2(0, \pi)} = 1. \quad (7)$$

Such systems can not form the unconditional basis. We show it more detailed.

The problem

$$-v''(x) = \bar{\lambda}v(x), \quad 0 < x < \pi; \quad (8)$$

$$v(0) + v(\pi) = 0, \quad v'(\pi) + \alpha v(\pi) = 0$$

is conjugated to the problem (1). The system of the eigenfunctions of this problem is biorthogonal to the system  $\{u_{k1}, u_{k2}\}_{k=0}^{\infty}$ :

$$\begin{aligned} v_{k1}(x) &= \frac{2}{\pi} \left\{ \sin((2k+1)x) - \frac{2k+1}{\alpha} \cos((2k+1)x) \right\} \\ v_{k2}(x) &= C_{k2} \left\{ \sin(2\beta_k x) - \frac{2\beta_k}{\alpha} \cos(2\beta_k x) \right\}, \quad k = 0, 1, 2, \dots \end{aligned} \quad (9)$$

The constant  $C_{k2}$  are taken from the biorthogonal relations  $(u_{k2}, v_{k2}) = 1$ . Since we will not use the explicit form of the biorthogonal system, then we do not present here the explicit form of constant  $C_{k2}$ .

Due to biorthogonality of the system, the equations

$$(u_{k1}, v_{k1}) = 1, \quad (u_{k2}, v_{k1}) = 0, \quad k = 0, 1, 2, \dots$$

are valid.

It follows that  $(u_{k1} - u_{k2}, v_{k1}) = 1$ . Using the Cauchy-Bunyakovsky inequality, we get the estimate from the bottom

$$\|v_{k1}\| \geq (\|u_{k1} - u_{k2}\|)^{-1}.$$

Since  $\|u_{k1}\| = \sqrt{\pi/2}$ , and  $\lim_{k \rightarrow \infty} \|u_{k2}\| = \sqrt{\pi/2}$ , then from here and from (7) it is easy to obtain

$$\lim_{k \rightarrow \infty} \|u_{k1}\| \|v_{k1}\| = \infty.$$

That is, the necessary condition of the basis property does not hold.

Lemma is proved.

It is necessary to note the fact, that the system of eigenfunctions  $\{u_{k1}, u_{k2}\}_{k=0}^{\infty}$  does not have the basis, also follows from more general facts [8].

### Forming the basis

Now from elements of the system  $\{u_{k1}, u_{k2}\}_{k=0}^{\infty}$  we construct a new system which will be a basis in  $L_2(0, \pi)$ . We introduce new functions

$$\begin{aligned} \varphi_{2k}(x) &= u_{k1}(x), \\ \varphi_{2k+1}(x) &= (u_{k2}(x) - u_{k1}(x)) (2\delta_k)^{-1}, \end{aligned} \quad k = 0, 1, 2, \dots \quad (10)$$

Let us show that the constructed system is a Riesz basis in  $L_2(0, \pi)$ .

The biorthogonal system to (10) has the form:

$$\begin{aligned} \psi_{2k}(x) &= v_{k2}(x) + v_{k1}(x), \\ \psi_{2k+1}(x) &= 2\delta_k v_{k2}(x), \quad k = 0, 1, 2, \dots \end{aligned}$$

This system is constructed from the eigenfunctions of the problem (8) conjugated to (1).

Let us show that the constructed additional system has the basis property.

**Lemma 2.** *The system of functions  $\{\varphi_k(x)\}_{k=0}^{\infty}$  forms a Riesz basis in  $L_2(0, \pi)$ .*

**Proof.** Since this system is constructed from the eigenfunctions of the problem with regular boundary conditions and with the help of non-degenerated linear combinations, then the completeness and minimality of the system do not change.

Let us prove asymptotic quadratic closeness of the system  $\{\varphi_k(x)\}_{k=0}^{\infty}$  to the system forming the Riesz basis. As such we choose the system of eigen- and associated functions of a problem of the Samarskii-Ionkin type:

$$\begin{aligned} -w''(x) &= \lambda w(x), \quad 0 < x < \pi; \\ w(0) &= 0, \quad w'(0) + w'(\pi) = 0. \end{aligned}$$

The boundary conditions of this problem are not strengthened regular. All the eigenvalues of this problem, except zero values, are multiple:  $\lambda_k^{(1)} = \lambda_k^{(2)} = (2k+1)^2$ ,  $k = 0, 1, 2, \dots$ . The eigenfunctions  $w_{2k}$  and the associated functions  $w_{2k+1}$  of the problem form the Riesz basis in  $L_2(0, \pi)$  and have the form:

$$w_{2k}(x) = \sin((2k+1)x), \quad k = 0, 1, 2, \dots; \quad w_{2k+1}(x) = x \cos((2k+1)x).$$

We need to show that the series converges

$$\sum_{k=0}^{\infty} \|\varphi_k - w_k\|^2 < \infty.$$

It is evident that  $\varphi_{2k} - w_{2k} = 0$ . For odd numbers we have:

$$\varphi_{2k+1}(x) = \frac{\sin(2\beta_k x) - \sin((2k+1)x)}{2\delta_k} = \frac{\sin(\delta_k x)}{\delta_k x} x \cos((2k+1 + \delta_k)x).$$

Thus it is not difficult to get the estimate  $|\varphi_{2k+1}(x) - w_{2k+1}(x)| \leq C\delta_k$ . From here and from the asymptotics (6) for  $\delta_k$  we have the asymptotic inequality  $|\varphi_{2k+1} - w_{2k+1}| \leq C_1/k$  where  $C_1$  does not depend on  $k$ .

The obtained inequality provides the quadratic closeness of the system  $\{\varphi_k(x)\}_{k=0}^{\infty}$  and the Riesz basis  $\{w_k(x)\}_{k=0}^{\infty}$ . Lemma is proved.

Further on, by standard methods it is not difficult to justify that if the function  $f(x) \in C^2[0, \pi]$  and satisfies the boundary conditions of the problem (1), then its Fourier series by the system  $\{\varphi_k(x)\}_{k=0}^{\infty}$  converges uniformly.

We can calculate that

$$\begin{aligned} -\varphi_{2k}''(x) &= \lambda_k^{(1)} \varphi_{2k}(x), \\ -\varphi_{2k+1}''(x) &= \lambda_k^{(2)} \varphi_{2k+1}(x) + \frac{\lambda_k^{(2)} - \lambda_k^{(1)}}{2\delta_k} \varphi_{2k}(x). \end{aligned} \tag{11}$$

Using these formulas, it is possible to apply the method of separation of variables for solving problems of the type (2) - (5).

### Use for solving of the nonlocal boundary equation

We can write any solution of problem (2) - (5) in the form of a biorthogonal series

$$u(r, \theta) = \sum_{k=0}^{\infty} R_k(r) \varphi_k(\theta), \tag{12}$$

where

$$R_k(r) = (u(r, \cdot), \psi_k(\cdot)) \equiv \int_0^\pi u(r, \theta) \psi_k(\theta) d\theta.$$

Functions (12) satisfy the boundary conditions (4) and (5).

Substituting (12) into equation (2) and the boundary conditions (3), taking into account (11), for finding unknown functions  $R_k(r)$  we obtain following problems:

$$\begin{aligned} r^2 R_{2k+1}''(r) + r R_{2k+1}'(r) - \lambda_k^{(2)} R_{2k+1}(r) &= 0, \\ r^2 R_{2k}''(r) + r R_{2k}'(r) - \lambda_k^{(1)} R_{2k}(r) &= \frac{\lambda_k^{(2)} - \lambda_k^{(1)}}{2\delta_k} R_{2k+1}(r), \end{aligned} \tag{13}$$

with the boundary conditions  $R_k(1) = f_k$ , where  $f_k$  are the Fourier coefficients of the expansion of the function  $f(\theta)$  into the biorthogonal series by  $\{\varphi_k(\theta)\}_{k=0}^{\infty}$ .

The regular solution of (13) exists, is unique and can be written in the explicit form:

$$\begin{aligned} R_{2k+1}(r) &= f_{2k+1} r \sqrt{\lambda_k^{(2)}}, \\ R_{2k}(r) &= f_{2k} r \sqrt{\lambda_k^{(1)}} + f_{2k+1} \frac{1}{2\delta_k} \left( r \sqrt{\lambda_k^{(2)}} - r \sqrt{\lambda_k^{(1)}} \right). \end{aligned} \quad (14)$$

Substituting (14) into (12), we obtain a formal solution of the problem:

$$\begin{aligned} u(r, \theta) &= \sum_{k=0}^{\infty} f_{2k} r^{2k+1} \sin((2k+1)\theta) + \\ &+ \sum_{k=0}^{\infty} f_{2k+1} \frac{1}{2\delta_k} [r^{2\beta_k} \sin(2\beta_k\theta) - r^{2k+1} \sin((2k+1)\theta)]. \end{aligned} \quad (15)$$

**Theorem** *If  $f(\theta) \in C^2[0, \pi]$ ,  $f(0) = 0$ ,  $f'(0) = -f'(\pi) + \alpha f(\pi)$ , then there exists a unique classical solution  $u(r, \theta) \in C^0(\bar{D}) \cap C^2(D)$  of the problem (2)-(5).*

**Proof.** The uniqueness of the classical solution of the problem follows from the maximum principle and the Zaremba-Giraud principle for the Laplace equation. The formal solution of the problem is shown in the form of (15). In order to make sure that these functions are really the desired solutions we need to verify the applicability of the superposition principle. For it we need to show the convergence of the series, the possibility of termwise differentiation, and to prove the continuity of these functions on the boundary of the half-disk.

The possibility of differentiating the series (15) any number of times at  $r < 1$  is an obvious consequence of the convergence of power series and two-sided estimates (6) for  $\delta_k$ . Let us justify the uniform convergence of the series (12) at  $r \leq 1$ . For this we use the sign of the uniform convergence of Weierstrass.

By direct calculation it is easy to see that the series (15) is majorized by the series  $C_1(|f_0| + |f_1| + |f_2| + \dots)$ . This series converges due to the requirements of the theorem imposed on  $f(\theta)$ . Since all the terms of the series (15) are continuous functions, then the function  $u(r, \theta)$  is continuous in the boundary domain  $\bar{D}$ .

The proof of the theorem is complete.

## Conclusion

Thus, in the present work we investigated a nonlocal boundary spectral problem (1) for an ordinary differential equation in an interval  $(0, \pi)$ . The boundary conditions of this problem are regular but not strengthened regular. The difference of this problem is: the system of eigenfunctions  $\{u_{k1}, u_{k2}\}_{k=0}^{\infty}$  of the problem (1) is complete and minimal, almost normed but does not form even an ordinary basis in  $L_2(0, \pi)$ .

Based on these eigenfunctions  $\{u_{k1}, u_{k2}\}_{k=0}^{\infty}$  there we constructed a special system of functions  $\{\varphi_k(x)\}_{k=0}^{\infty}$  that already forms a Riesz basis in  $L_2(0, \pi)$ .

This fact is used for solving of the nonlocal boundary equation (2) - (5).

## Acknowledgements

The authors express their gratitude to T.Sh. Kalmenov and M.A. Sadybekov for valuable advices during the work.

This work was supported by the grant 0824/GF4 of the Ministry of Education and Science of Republic of Kazakhstan.



### References

- [1] *Il'in V.A.; Kritskov L.V.* Properties of spectral expansions corresponding to non-self-adjoint differential operators // Journal of Mathematical Sciences. Vol.116, No.5, p.3489-3550, (2003)
- [2] *Locker J.* Spectral Theory of Non-Self-Adjoint Two-Point Differential Operators. V.192 of Mathematical Surveys and Monographs. Amsterdam: North-Holland, 2003.
- [3] *Makin A.S.* On Summability of Spectral Expansions Corresponding to the Sturm-Liouville Operator // International Journal of Mathematics and Mathematical Sciences. Vol.2012, Article ID 843562, p.1-13, doi:10.1155/2012/843562, (2012)
- [4] *Naimark M.A.* Linear Differential Operators. New York: Ungar, 1967.
- [5] *Mihailov V.P.* On Riesz bases in  $L_2(0, 1)$  // Doklady Akademii Nauk SSSR. Vol.144, No.5, p.981-984, (1962) (in Russian)
- [6] *Kesel'man G.M.* On the unconditional convergence of eigenfunction expansions of certain differential operators // Izv. Vuzov. Mat.. Vol.2, No.39, p.82-93, (1964) (in Russian)
- [7] *Dunford N.; Schwartz J.T.* Linear Operators, Part III. New York: John Wiley & Sons, 1971.
- [8] *Lang P.; Locker J.* Spectral theory of two-point differential operators determined by  $D^2$ . II. Analysis of case // Journal of Mathematical Analysis and Applications. Vol.146, No.1, p.148-191, (1990)
- [9] *Moiseev E.I.; Ambartsumyan V.E.* On the solvability of nonlocal boundary value problem with the equality of flows at the part of the boundary and conjugated to its problem // Differential Equations. Vol.46, No.5, p.718-725, (2010)
- [10] *Moiseev E.I.; Ambartsumyan V.E.* On the solvability of nonlocal boundary value problem with the equality of flows at the part of the boundary and conjugated to its problem // Differential Equations. Vol.46, No.6, p.892-895, (2010)
- [11] *Mokin A.Yu.* On a family of initial-boundary value problems for the heat equation // Differential Equations. Vol.45, No.1, p.126-141, (2009)

*Поступила в редакцию 27 февраля 2015 года*